

Fundamentals of Glycan Structure

Learning Objectives

- ☐ *How are glycans named?*
- ☐ *What are the different constituents of a glycan?*
- ☐ *How are these represented?*
- ☐ *What conformations do sugar residues adopt in solution*
- ☐ *Why do glycan conformations matter?*

Fundamentals of Glycan Structure

- ❑ ***Carbohydrate Nomenclature***
- ❑ ***Monosaccharides***
 - ✓ *Structure*
 - ✓ *Fisher Representation*
 - ✓ *Cyclic Form*
 - ✓ *Chair Form*
 - ✓ *Mutarotation*
- ❑ ***Monosaccharide Derivatives***
 - ✓ *Reducing Sugars*
 - ✓ *Uronic Acids*
 - ✓ *Other Derivatives*
- ❑ ***Monosaccharide Conformation***
- ❑ ***Inter-Glycosidic Bond***
 - ✓ *Normal*
 - ✓ *Sucrose*
 - ✓ *Lactose*
 - ✓ *Sequence Specificity and Recognition*
 - ✓ *Branching*



Carbohydrate Nomenclature

- ❑ **The word 'carbohydrate' implies "hydrate of carbon" ... $C_n(H_2O)_m$**
 - ✓ Glucose (a monosaccharide) $C_6H_{12}O_6$... $C_6(H_2O)_6$
 - ✓ Sucrose (a disaccharide) $C_{12}H_{22}O_{11}$... $C_{12}(H_2O)_{11}$
 - ✓ Cellulose (a polysaccharide) $(C_6H_{12}O_6)_n$... $(C_6(H_2O)_6)_n$
- ❑ **Not all carbohydrates have this formula ... some have nitrogen**
 - ✓ Glucosamine (glucose + amine) ... $C_6H_{13}O_5N$... $-NH_2$ at the 2-position of glucose
 - ✓ N-acetyl galactosamine (galactose + amine + acetyl group) ... $C_8H_{15}O_6N$... $-NHCOCH_3$ at the 2-position of galactose
- ❑ **Typical prefixes and suffixes used in naming carbohydrates**
 - ✓ Suffix = '**-ose**' & prefix = '**tri-**', '**tetr-**', '**pent-**', '**hex-**'
 - ✓ Pentose (a five carbon monosaccharide) or hexose (a six carbon monosaccharide)
- ❑ **Functional group types**
 - ✓ Monosaccharides with an aldehyde group are called aldoses ... e.g., glyceraldehyde
 - ✓ Those with a keto group are called ketoses ... e.g., dihydroxyacetone

Monosaccharides


□ **Structure**

- ✓ *Have a general formula $C_nH_{2n}O_n$ and contain a carbonyl group*
- ✓ *Common monosaccharides have 3 \rightarrow 9 carbons*
- ✓ *Two molecules form the class of smallest monosaccharides ...
glyceraldehyde and dihydroxyacetone*





Monosaccharides

- *Glyceraldehyde has a chiral carbon*







* = chiral carbon
... 2 stereoisomers

- *3-Dimensional arrangement*





- *2-Dimensional representation*



Monosaccharides

□ *Emil Fischer's representation*














- ✓ *Arbitrarily assigned the dextrorotatory enantiomer as D-glyceraldehyde, which fortuitously proved correct*



Note: *This does not mean that all D-saccharides are dextrorotatory; likewise it does not mean that all D-saccharides are also (R) in the Cahn-Ingold-Prelog system*

Monosaccharides


☐ *'D' monosaccharides*




- ✓ *'D' monosaccharides have the same configuration on their penultimate carbon as 'D' glyceraldehyde*
- ✓ *Exactly similar series for 'L' monosaccharides starting with 'L' glyceraldehyde*
- ✓ *Likewise, similar series exists for 'L' and 'D' ketoses starting from dihydroxyacetone*

Monosaccharides


☐ 'L' ketoses




Dihydroxyacetone
(no D- or L-form)




L-erythrulose




L-xylulose




L-ribulose




L-psicose



L-fructose



L-sorbose



L-tagatose

Monosaccharides


❑ *Some Useful Information on Monosaccharides*

- ✓ *D-glucose is also called dextrose because it is dextrorotatory.*
- ✓ *D-glucose is the most abundant monosaccharide; it is present in most fruit juices*
- ✓ *D-glucose is also called blood sugar it is present in about 65 – 100 mg per 100 mL of blood*
- ✓ *The dextrose solution for iv drip contains about 5% glucose + 0.15% saline (KCl)*
- ✓ *Fructose is one of the monosaccharides present in disaccharide sucrose (table sugar) ... the other is D-glucose*
- ✓ *Galactose is part of lactose (milk sugar)*

Monosaccharides

❑ *Monosaccharides Are Typically not Linear (or Open-Chain)*

- ✓ Remember from organic chemistry: Aldehydes and ketones react with alcohols to give **hemi-acetals**




- ✓ What is an anomer?
- ✓ How are α - and β -anomers defined?
- ✓ What is an epimer?


Monosaccharides

❑ *Cyclic Structure of Ketoses*

- ✓ *Similar cyclic structures exist for five membered saccharides ... e.g., ribose.*
- ✓ *These five membered cyclic hemiacetals are well known for nucleic acids ... e.g., RNA (**ribo**nucleic acid) and DNA (**deoxyribo**nucleic acid)*







HAWORTH Projections



Monosaccharides

❑ *The Cyclic Form Assumes Different Conformations in Solution*




Chair Conformations

✓ *Why is β -D-glucose the most common sugar in nature?*

Monosaccharides


❑ *The Cyclic Form of D-Galactose*



Chair Conformations

Monosaccharides

❑ *Mutarotation Arises From Cyclic and Open-Chain Form Interconversion*



Optical activity

$$[\alpha]_D = +112.2^\circ$$


$$[\alpha]_D = +18.7^\circ$$

At equilibrium $[\alpha]_D = +52.7^\circ$ (Not an average!

But weighted average; 63.6% of the β-anomer)

Monosaccharide Derivatives


❑ *Reducing Sugars and Reducing End*



- ✓ *Reducing sugars and diabetes*
 - Hemoglobin (Hb) is glycated HbA1c levels should be < 6.5%
 - Reaction of high glucose with NH_2 group at the N-terminus (Val) of Hb
 - Non-enzymatic
 - Reflects Glc levels for the past 3 months (typical life of erythrocytes)

Monosaccharide Derivatives

□ *Uronic Acids*







✓ *Biological relevance*

- Several other uronic acids occur in our body, e.g., D-mannuronic acid (from D-mannose), L-iduronic acid (from L-idose), D-galacturonic acid
- Components of many polysaccharides
- Add to the conformational complexity, e.g., L-iduronic acid can exist in several conformations other than chair – ²S_O, ⁰S₂, ¹C₄ and ⁴C₁

Monosaccharide Derivatives

☐ **Glycosamines**

- ✓ Several aminosugars are known including D-glucosamine, D-mannosamine, D-galactosamine and N-acetyl-D-glucosamine. These have a nitrogen at position 2




☐ **Other Derivatives**

- ✓ **Glycan Esters**
 - The –OH groups can be esterified enzymatically, e.g., phosphate esters, acyl esters, and sulfate esters
- ✓ **Deoxy Sugars**
 - The –OH group can be replaced with –H, e.g., 2-deoxyribose (C-2 deoxy), fucose (C-6 deoxy), etc.
- ✓ **Methylated Sugars**
 - The –OH group can be methylated (methyltransferases), e.g., –OMe group at C-1

Monosaccharide Conformation










□ *Finding the Most Stable Conformation of a Monosaccharide*



D-Glucose

Monosaccharide Conformation


□ Conformation of L-Iduronic Acid




Generation of Oligosaccharides

❑ *The Inter-Glycosidic Bond*

- ✓ *The bond between two monosaccharides is the inter-glycosidic bond*
- ✓ *Fundamental linkage for generation of oligosaccharides*
- ✓ *Any hydroxylated molecule (e.g., amino acids Tyr, Ser, Thr; OR alcohols cholesterol, cholic acid; OR saccharides) can form glycosidic bond*
- ✓ *Most oligosaccharides form glycosidic bond between the anomeric carbon (C-1) and other carbons (C-2, C-3, C-4, and C-6).*
- ✓ *These different glycosidic linkages enhance the diversity of glycans*
- ✓ *Glycosidic bond induces directionality in the saccharide chain, i.e., non-reducing end → reducing end*
- ✓ *Glycosidic bonds make the oligomer flexible*



Glc α 1-4Glc
(maltose)




Glc β 1-6Glc
(gentiobiose)

Generation of Oligosaccharides

❑ *Sucrose's Inter-Glycosidic Bond is Different*

- ✓ *Sucrose (table sugar) is a disaccharide made from α -D-glucose and β -D-fructose, where the inter-glycosidic bond is α -1,2*




- ✓ *How do we find out which atoms are connected in polysaccharides and what is their orientation?*
- ✓ *How do we know what conformation does a sugar adopt?*

Generation of Oligosaccharides

❑ Lactose

- ✓ Lactose (milk sugar) is a disaccharide made from β -D-galactose and D-glucose, where the inter-glycosidic bond is β -1,4




- ✓ Lactose intolerance? ... cow's milk ... 4 – 6% lactose ... some infants, most blacks and many Orientals have low level of lactase, which hydrolyzes the β (1,4) bond of lactose ... lactose accumulates in colon where bacterial fermentation produces large quantities of CO_2 , H_2 and organic acids

Generation of Oligosaccharides

❑ *Sequence Specificity and Recognition*

- ✓ *Is lactose (D-galactose- β (1 \rightarrow 4)-D-glucose) is same as D-glucose- β (1 \rightarrow 4)-D-galactose?*




- ✓ *Remember! D-glucose is consumed by our body; L-glucose is left untouched!*

Generation of Oligosaccharides


❑ **Branching is a Unique Structure Only Present in Carbohydrates**

- ✓ Two inter-glycosidic linkages to the same monosaccharide results in branching
- ✓ Branching is not found in proteins, nucleic acids and lipids Only in carbohydrate



Neu5Ac α 6Gal β 4GlcNAc β 2Man α ₆
Man β 4GlcNAc β 4GlcNAc β -Asn
Neu5Ac α 6Gal β 4GlcNAc β 2Man α ₃

N-Glycan



Fuc α ₃
Neu5Ac α 3Gal β 4GlcNAc β ₆
Neu5Ac α 3Gal β 3GalNAc α -Ser

O-Glycan

Taken from 'Essentials of Glycobiology.' Edited by Varki, A.; et al.; Cold Spring Harbor Laboratory Press, New York, 2009, pg. 36.